

Name:

PID:

- 
- Print your *NAME* on every page and write your PID in the space provided above.
  - Show all of your work in the spaces provided. No credit will be given for unsupported answers, even if correct.
  - Supporting work for a problem must be on the page containing that problem. No scratch paper will be accepted.
  - No calculators, tables, phones, or other electronic devices are allowed during this exam. You may use your double-sided handwritten notes, but no books or other assistance.
- 

**DO NOT TURN PAGE UNTIL INSTRUCTED TO DO SO**  
(This exam is worth 25 points)

**Problem 0.**(1 point.) Follows the instructions on this exam and any additional instructions given during the exam.

**Name:**

**Problem 1.**(6 points.) Suppose we have a standard six-sided dice, and we roll it 100 times. Let  $X_i$  be the number on the face of the dice for roll  $i$ . Let  $X$  be the sum of the dice rolls, i.e.,  $X = \sum_{i=1}^n X_i$ .

- a) (3 points) Find  $\mathbb{E}[X_i]$ ,  $\mathbb{E}[X]$ , and  $\text{Var}[X]$ .
- b) (3 points) Use Chebyshev's inequality to bound  $\mathbb{P}(|X - 350| \geq 60)$ .

**Name:**

**Problem 2.**(6 points.) Let  $X_1, \dots, X_n$  be independent random variables where

$$X_i = \begin{cases} 1, & \text{with probability } 1/3 \\ 0, & \text{with probability } 1/3 \\ -1, & \text{with probability } 1/3. \end{cases}$$

Let  $X = \sum_{i=1}^n X_i$ . Bound  $\mathbb{P}(|X - \mathbb{E}[X]| \geq \sqrt{n})$  by using the following inequalities.

- a) (3 points) Chebyshev's inequality.
- b) (3 points) Chernoff's inequality.

**Name:**

**Problem 3.**(6 points.)

- a) (2 points) Given  $\vec{u}$  and  $\vec{v}$  in  $\mathbb{R}^n$ , show that if  $A = \vec{u}\vec{v}^T$ , then  $\|A\|_2 = \|\vec{u}\|_2\|\vec{v}\|_2$ .
- b) (4 points) Given  $\vec{u} = (1, -1, 1)^T$  and let  $M = \vec{u}\vec{u}^T$ . Find 2-norm and 1-norm of  $M$ .  
(Hint: you can use Part a)

**Name:**

**Problem 4.**(6 points.) Given an  $n \times m$  matrix  $A$  with entries in  $\{0, 1\}$  and a vector  $\vec{b} \in \mathbb{R}^m$ , i.e.,

$$A = [a_{ij}] = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1m} \\ a_{21} & a_{22} & \cdots & a_{2m} \\ \vdots & \vdots & \cdots & \vdots \\ a_{n1} & a_{n2} & \cdots & a_{nm} \end{bmatrix}, \text{ and } \vec{b} = \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_m \end{bmatrix}.$$

- a) (3 points) Write a formula for  $\|A\vec{b}\|_\infty$ .
- b) (3 points) Suppose that  $b_j$ 's are independent random variables, where  $b_j = -1$  with probability  $1/2$  and  $1$  else. Let  $Z_1 = \sum_{j=1}^m a_{1j}b_j$ . Use Chernoff's bound inequality to bound

$$\mathbb{P}(|Z_1| \geq \sqrt{4m \ln n})$$